

SUPREX™ K29

Foliar Potassium Fertilizer

0-0-29

GUARANTEED ANALYSIS

Soluble Potash (K₂O).....29%

Derived from potassium acetate

GENERAL INFORMATION

Suprex™ K-29 is a highly available liquid formulation of potassium acetate. Suprex K-29'S low salt index and superior foliar absorption quickly increases potassium levels in plants with minimal potential for burn, even at higher rates. Suprex K-29'S organic nature provides greater absorption of potassium than other commonly used potassium sources and is safe for plants even at higher rates. Suprex K-29 is especially effective when potassium utilization is greatest, such as during rapid growth periods, fruit development and fruit sizing.

KEEP OUT OF REACH OF CHILDREN



CAUTION

READ ENTIRE LABEL BEFORE USING

PRECAUTIONARY STATEMENTS:

Avoid prolonged or repeated contact with eyes, skin, and clothing. Chemical goggles or a full face shield should be worn. To protect skin wear appropriate protective equipment, such as rubber or plastic aprons, rubber gloves, and boots. Avoid breathing mist or vapor. Keep containers closed. Wash thoroughly after handling. May cause gastrointestinal distress if swallowed.

F1129

APPLICATION INSTRUCTIONS

Foliar Applications:

Turfgrass and Sod (including greens, fairways and lawns): Apply 2 to 8 quarts/acre every 14 to 21 days as needed. Greens and Tees: Apply 2 to 8 ounces per 1000 sq. ft. every 14 to 21 days as needed. Corn: Apply 2 to 6 quarts/acre prior to tasseling and repeat in 7 to 10 days.

Ornamental Plants and Trees: Apply 2 to 6 quarts/100 gallons of water foliarly every 14 to 21 days as needed. Apply 6 quarts/100 gallons of water as a drench or in transplant water.

Cotton: Apply 0.5 to 2 gallons/acre at early bloom repeating up to 4 applications at 7-10 day intervals.

Fruits, Nuts and Vine Crops (Apples, Citrus, Grapes, Peaches, etc.): Apply 2 to 8 quarts/acre in 100 gallons of water beginning at first full leaf and/or fruit expansion, repeat as needed.

Grains: Apply 2 to 6 quarts/acre at tillering to early boot stage.

Legumes: Apply 2 to 6 quarts/acre at late bud to 10% bloom.

Rice: Apply 0.5 to 2 gallons/acre at panicle initiation.

Small fruits: (Blackberries, Blueberries, and Strawberries) Apply 2 to 6 quarts/acre every 10 to 14 days starting before bloom.

Tuber and Root Crops: Apply 2 to 3 gallons/acre at tuber initiation with second application 2 to 3 weeks later and final application at tuber bulking.

Vegetables, Cucurbits, Fruiting and Brassicas: Apply 2 to 8 quarts/acre starting at fruit formation or bottom establishment. Apply every 7 to 14 days as needed.

Vegetables, Leafy: Apply 1 to 6 quarts/acre beginning at thinning.

Aerial Applications:

Use 2 to 5 quarts/acre in 3 to 10 gallons of water of spray solution.

Sprinkler/Drip Irrigation:

Tree - under canopy: Use 5 to 10 gallons/acre per application beginning at full leaf and schedule as necessary every 10 to 14 days.

Vegetable, vines and row crops: Use 3 to 6 gallons/acre per application, make sequential applications based on crop development every 10 to 14 days. Higher rates are for Flood and Furrow Irrigation.

Pivot/Overhead Irrigation:

Vegetable and row crops: Apply 2 to 4 gallons/acre per application as needed every 10 to 14 days. Trees: Apply 2 to 10 gallons/acre. After injection, allow irrigation to run long enough to wash residual fertilizer from foliage.

Sprinkler/Drip Irrigation:

Tree - under canopy: Use 5 to 10 gallons/acre per application beginning at full leaf and schedule as necessary every 10 to 14 days.

Vegetable, vines and row crops: Use 3 to 6 gallons/acre per application, make sequential applications based on crop development every 10 to 14 days. Higher rates are for Flood and Furrow Irrigation.

Pivot/Overhead Irrigation:

Vegetable and row crops: Apply 2 to 4 gallons/acre per application as needed every 10 to 14 days. Trees: Apply 2 to 10 gallons/acre. After injection, allow irrigation to run long enough to wash residual fertilizer from foliage.

MIXING INSTRUCTIONS

Use caution when tank mix pH goes below a pH of 6. A jar test is recommended to test compatibility when new or unfamiliar tank mix combinations are used. When used as directed this product does not supply all the nutrients required by plants and is designed to supplement a soil fertility plan.

STORAGE AND DISPOSAL

STORAGE: Store at ambient temperatures in dry secure area. Keep container tightly closed. Do not allow water to be introduced into the container contents.

DISPOSAL: Dispose of empty containers by triple rinsing or equivalent and discard empty containers in accordance with current Local, State and Federal regulations. Do not contaminate water, food or feed sources by cleaning of equipment or disposal of waste or rinsate.

CONDITIONS OF SALE

Seller warrants this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions and use. No one other than an officer of the seller is authorized to make any warranty, guarantee, or direction concerning this product. Because time, place, rate of application and other conditions of use are beyond the sellers control, the sellers liability from handling, storage, and use of this product is limited to replacement of the product or a refund of the purchase price.

Do not apply if a heavy rain is expected. Do not apply near water, storm drains or drainage ditches or to any impervious surface. Only apply to the intended application area.

PERFORMANCE
NUTRITION®

Guaranteed and Distributed By:
LidoChem LLC - Performance Nutrition Division

20 Village Court, Hazlet, NJ 07730

Phone: (732) 888-8000 | Email: info@pnfertilizers.com

Performance Nutrition and Suprex are trademarks of LidoChem LLC ©2025 LidoChem LLC

Weight per Gallon: 11.1 lb. (5.0 kg.)

Net Volume and Weight

<input type="checkbox"/>	5 Gal. (19 Lit)	55.5 lb. (25.2 kg.)
<input type="checkbox"/>	55 Gal. (208 Lit)	610.5 lb. (277 kg.)
<input type="checkbox"/>	250 Gal. (945 Lit)	2,775 lb. (1,259 kg.)